
ANALYTICAL STUDY OF THE LEADING CAUSES OF DEATH OF PALLIATIVE PATIENTS

Smiiianov V.A.1, Hubert-Lutecka A.²

¹Sumy State University, Sumy, Ukraine

²University of Rzeszow, Rzeszow, Poland

<https://doi.org/10.35339/ic.11.4.smh>

ABSTRACT

Background. Palliative and Hospice Care (PHC) aims to prevent premature death of patients from complications of serious illnesses and their comorbid impact on vital functions. As well as adequate pain relief, treatment should be aimed at alleviating suffering and improving the quality of life of such patients. The causes of death of palliative patients, which are closely related to the PHC organization, in particular to the package budget financing of inpatient and mobile palliative care for adults and children, the leading needs of palliative patients, remain insufficiently studied.

Aim. Analysis of the main causes of death of palliative patients depending on the diagnosis and determination of their impact on the organization of palliative and hospice care.

Materials and Methods. The method of system analysis, comparative method and bibliosemantic method were used for the research.

Results and Conclusions. Causes of death were studied for diseases listed as palliative in severe cases and in the presence of complications. The analysis allows us to deepen our understanding of the practical aspects of organizing palliative and hospice care in meeting the needs of palliative patients, adults and children. The summary of the causes of death allows us to group these causes into organ and system failure; vascular crises; asphyxia; thromboembolic conditions; infectious complications up to sepsis; chronic intoxication; tumor growth in other organs with disruption of their vital functions, metastasis; malignancy of benign tumors; complications of treatment; brain and spinal cord lesions due to epileptic seizures, injuries, inflammatory processes; endocrine comas; gangrene and bedsores; severe immunodeficiency; underdevelopment or absence of organs in congenital malformations; prematurity; suicides in depressive states.

Keywords: *palliative and hospice care, primary palliative diagnosis, comorbidity.*

Introduction

Palliative and Hospice Care (PHC) is provided to terminally ill patients in the last days of their lives and is intended to reduce their suffering, the feeling of pain and to prepare for a dignified death. Modern development of medical technologies theoretically allows to ensure painless dying, which is possible with the correct organization of medical care. The countries with developed PHC systems offer patients recognition of their palliative status, stay in specialized institutions (hospices, palliative departments and wards) or in "hospices at home", adequate pain relief, euthanasia as an op-

tion, as well as social support and spiritual support, which extend to both the patient himself and his loved ones (caregivers) [1–3].

A significant part of palliative diseases belongs to the category of disabling and is accompanied by chronic intense pain [4; 5]. Depending on the main palliative diagnosis and comorbid pathology, in severe palliative conditions, nociceptive (associated with nerve irritation), neuropathic (associated with organic nerve damage) type of pain prevails, or a combination of these types of pain is present. Depending on the intensity of pain, which is derived from the characteristics of the pathological process and individual perception of pain, narcotic drugs (morphine, oxycodone, buprenorphine, etc.), non-narcotic drugs (primarily Nonsteroidal Anti-Inflammatory Drugs, NSAIDs) and adjuvants (not intended for direct pain relief, but capable of enhancing the effect of analgesic drugs – sedatives, hypnotics, anticonvulsants, neuroleptics, etc.)

Corresponding Author:

Smiiianov Vladyslav A. – MD, DMedSc, Professor,
Head of the Public Health Department of the Sumy
State University Medical Institute, Ukraine.

✉ Ukraine, 40004, Sumy, Pratsi str., 3.

E-mail: v.smiiianov@med.sumdu.edu.ua

are used for pain relief [2; 6; 7]. The recent legalization of medical cannabis theoretically significantly expands the possibilities of adequate pain relief. However, to practically realize this opportunity, Ukraine needs to accelerate the adoption of by-laws regulating cultivation of cannabis, production of cannabis-based drugs, and simplify the procedures for obtaining these drugs by palliative patients [8; 9].

Pain in severe palliative conditions is not a "signal" pain, i.e., it does not indicate a previously unknown health problem. On the contrary, it is emotionally debilitating and depletes the immune and endocrine systems. Its impact on inflammatory processes is negative due to its participation in "pathological circles" [2; 4; 5; 10]. This indicates the need for constant adequate pain relief for palliative patients, but taking into account the risks of excessive sedation, which can lead to death. It is these risks that allow us to talk about the need to prevent premature death in patients who already expect it to be imminent. In the ongoing debate about the risks and benefits of long-term pain relief with narcotic drugs, it is necessary to focus on evidence collected according to the rules of evidence-based medicine [11; 12].

In recent years, researchers from the Ukrainian Center for Public Data (Kyiv, Ukraine) and Kharkiv National Medical University have made progress in determining the list of palliative diseases in Ukraine [13–16]. Representatives of the first organization calculated the need for PHC and determined the list of main palliative diagnoses of adults and children for 2018. Representatives of the second organization expanded this list, calculated the need for PHC for 2019–2020 and forecasted this need for 2021–2022. Subsequently, this forecast was checked and revised retrospectively using the creeping trend method with a constant smoothing segment. The method itself was also improved [16; 17].

In their further studies, representatives of the Kharkiv group studied the features of the organization of PHC at the regional level [18; 19], improved the questionnaire on the quality of life of palliative patients of neurological and oncological profiles [20; 21], studied the issues of types of pain in different palliative diagnoses, features of pain relief [4; 6; 7], attitudes towards the possibility of legalizing euthanasia [22–25]. To date, the causes of death of palliative patients, which are closely related to the organization of PHC, in particular, with the package budget financing of stationary and mobile palliative care for adults and

children [26], and the leading needs of palliative patients [1; 8], remain insufficiently studied in the works of the authors and according to the results of searches on PubMed, Google and Google Scholar.

The **aim** of the study was to analyze the main causes of death of palliative patients depending on the diagnosis and to determine their impact on the organization of palliative and hospice care.

Materials and Methods

The research used methods of systematic analysis, comparative and bibliosemantic, with a search for sources on PubMed, Google and Google Scholar using the keywords "palliative and hospice care", "causes of death of palliative patients", "comorbidity of palliative diseases" in Ukrainian and English. In connection with the theoretical approach to the research, bioethical examinations of the research materials were not conducted. Statistical methods were also not used.

Results and Discussion

The causes of death of cancer patients (C00–C97, D00–D48 according to the International Statistical Classification of Diseases and Related Health Problems, 10th revision, ICD-10) are cancer intoxication, failure of affected organs and systems (respiratory, cardiovascular, renal, hepatic failure), cachexia, metastasis and germination of tumors to other organs with disruption of their vital functions, thromboembolism, infectious complications (sepsis, infection of decaying tumors). Treatment complications are also possible. For example, death due to excessive sedation with narcotic analgesics [7; 27; 28]. Cancer patients with malignant neoplasms accounted for (36.3–40.4)% of all palliative adult patients and (13.5–17.8)% of children with palliative diagnoses in Ukraine for the period 2018–2020, according to the calculation of the need for PHC in Ukraine performed by Nesterenko V.G. et al. (2021) [14–17].

The causes of death from cardiovascular diseases (I00–I99 according to ICD-10) are usually acute and chronic heart failure, circulatory failure due to myocardial infarction and stroke of the brain vessels (hemorrhagic and ischemic), chronic heart failure, thromboembolic conditions and cardiac arrhythmias. Rupture of aneurysms against a background of hypertension with internal bleeding is also possible [29–33]. Cardiovascular diseases accounted for (32.8–40.1) % of all palliative adult patients and (1.6–2.3) % of children with palliative diagnoses in Ukraine for the period 2018–2020 according to the calculation of the need for PHC in Ukraine.

The causes of death in adult dementia (F72–F79 according to ICD-10) and severe and profound mental retardation in children (F00–F03 according to ICD-10) are swallowing disorders and related asphyxia and aspiration pneumonia; urinary tract infections up to septic conditions; pressure ulcers and their infectious complications; anorexia; treatment complications [34–37]. An important aspect of organizing PHC for these categories of palliative patients is the resumption of the collection of statistical data on morbidity, which was discontinued in Ukraine in 2019.

The causes of death from tuberculosis (A15–A19 according to ICD-10) are bronchial obstruction and respiratory failure; massive pulmonary hemorrhage; HIV infection (in combination with tuberculosis, the stage of this disease is defined as AIDS); meningitis. The infectious process threatens the life of the patient when infected with a pathogen with multidrug resistance to antibiotics [38–41]. In 2022, tuberculosis was the second leading cause of death from infectious diseases in the world, second only to COVID-19 (7.5 million new cases of infection). In 2023, it was in first place (8.2 million new cases of infection). At the same time, the elderly and people with disabilities are at risk and often require palliative care [42–45].

The causes of death in diabetes mellitus (E10–E14 according to ICD-10) are diabetic coma, vascular crises, renal failure, infectious complications (up to sepsis) due to the development of gangrene of the lower extremities, and pressure sores [46].

The causes of death in rheumatoid arthritis (M05–M06 according to ICD-10) are mainly infectious complications (up to sepsis) of immobility caused by the disease (bedsores, pneumonia) and complications of treatment (primarily immunosuppressants, which reduce the body's resistance to infectious diseases and can cause liver fibrosis) [47–49]. Fibrosis and cirrhosis of the liver (K74 according to ICD-10) are also considered as palliative conditions in adults and children. Most often, their occurrence is associated with chronic hepatitis (K73, K75.2, K75.3 according to ICD-10), and death is caused by liver failure, bleeding, malignancy (liver cancer), infectious complications of immobility in these diseases (skin infections, pneumonia), which can cause sepsis [50].

The causes of death in HIV infection are the development of the final stage of the disease (AIDS) and related infectious and oncological processes (B20–B24 according to ICD-10). Due to the ineffectiveness of immune defense, even

opportunistic infections can cause severe consequences, and the range of possible oncological diagnoses is significantly expanded (for example, due to Kaposi's sarcoma). NeuroAIDS can also be a direct cause of death. In war conditions, temporarily displaced persons and migrants with HIV-positive status may be deprived of access to antiretroviral therapy, which can accelerate the development of AIDS and its fatal complications [51–53].

The causes of death in Chronic Obstructive Pulmonary Disease (COPD) (J43–J47 according to ICD-10) are respiratory failure due to pneumonia, bronchial obstruction, destruction of lung tissue affected by the infectious process, and tumor destruction. Treatment complications are also possible: allergic reactions up to anaphylaxis, cardiac arrhythmias, thromboembolism, steroid insufficiency due to long-term treatment with inhaled corticosteroids. Chronic lack of oxygen can cause heart failure. The onset of death is significantly accelerated by the widespread use of tobacco smoking in the world [54–57]. Even before the start of the COVID-19 pandemic, COPD was one of the leading causes of death in the world. Thus, in 2016, the disease caused more than 3 million deaths and took 3rd place among the 10 leading causes of death [56]. When collecting data on causes of death, COPD should be counted separately from deaths caused by COVID-19. But in practice, these causes were often not separated.

The causes of death in kidney diseases (N00–N15, N20–N23 according to ICD-10) are renal failure, uremia, infectious complications of inflammatory processes in the kidneys. Severe complications of treatment are also possible: the development of an infectious process at the site of access to the vessels during dialysis, malignancy of kidney tumors during dialysis, gastrointestinal bleeding from damage to the mucous membranes by oral drugs, cardiac arrhythmias due to deviations from the norm of potassium content in the blood caused by treatment with loop and thiazide diuretics, angiotensin-converting enzyme inhibitors [58–63].

The causes of death in cystic fibrosis (E84 according to ICD-10) are respiratory, cardiovascular and hepatic failure, cystic fibrosis of the pancreas and infectious complications of the disease, up to sepsis [64; 65]. Insufficiency of organs and organ systems is also a cause of death in children with mucopolysaccharidosis (E76 according to ICD-10), but in addition to cardiovascular, respiratory and hepatic failure, renal failure should be added.

Other causes of death are seizures that damage the brain, arrhythmias and infectious complications [66; 67].

The causes of death in congenital malformations of newborns (Q00–Q99 according to ICD-10) are the absence or underdevelopment of organs, causing all possible types of organ failure, immediately after birth or during the growth and development of the child. Malformations that are present in the nervous system can cause seizures and paralysis. Complications of surgical treatment of these patients can also affect vital signs [68–71].

Perinatal conditions that lead to death and require palliative care for children (P05–P96 according to ICD-10) may be associated with congenital malformations [72], prematurity, pathology of pregnant women (including extragenital), fetal asphyxia, bleeding of pregnant women and women in labor, infections of pregnant women, fetus and newborn [73–75].

The causes of death in cerebral palsy (G80 according to ICD-10) may include decompensation of the underlying disease with impaired nervous regulation of the respiratory process and the development of acute heart failure, aspiration pneumonia, thromboembolic conditions, seizures, tumor malignancy, infectious processes of the urinary system up to sepsis [76–78].

Inflammatory diseases of the Central Nervous System (CNS) (E70.0 according to ICD-10) lead to death due to depression of the respiratory center, additional damage to nerve structures during epileptic seizures, as well as suicides in a state of depression, malignancy of tumors and complications of treatment [79–81]. In turn, epilepsy (G40 according to ICD-10) can cause Sudden Unexpected Death in Epilepsy (SUDEP) due to damage to nerve centers during an attack with a stroke, trauma (especially when falling during an attack). Drowning during an attack and suicide in depressive states are also possible [82; 83].

Multiple sclerosis (G35 according to ICD-10) can cause death due to the development of pneumonia, vascular crises, pulmonary embolism, infectious complications of immobility (bedsores, urinary tract infections), up to sepsis, and suicide in depressive states [84–87].

The medical and social component of preventing most premature deaths of palliative patients consists of promoting timely examination and treatment of diseases with controllable and condi-

tionally controllable risk factors, promoting vaccinations, safe sex, quitting tobacco smoking, recreational drug use and alcohol abuse, implementing anti-epidemic rules, and building partnerships between patients and doctors in health care [88–93].

Conclusions

The causes of death have been studied for diseases listed as palliative in severe cases and in the presence of complications. The analysis allows us to deepen our understanding of the practical aspects of organizing palliative and hospice care in terms of meeting the needs of palliative patients, adults and children. The summary of determining the causes of death allows us to group these causes into organ and system failure (cardiovascular, respiratory, hepatic and renal); vascular crises (cerebral strokes and myocardial infarctions, cardiac arrest due to rhythm disturbances); asphyxia (fetal, adults due to food aspiration, bronchial obstruction); thromboembolic conditions (including pulmonary embolism); infectious complications up to sepsis, chronic intoxication (cancerous, including during tumor decay; uremia; against the background of multi-resistance of microflora to antibiotics); tumor growth in other organs with disruption of their vital functions, metastasis; malignancy of benign tumors; complications of treatment (including excessive sedation, which caused premature death; electrolyte imbalance; anaphylaxis of unsuccessful surgical interventions and resuscitation); brain and spinal cord lesions due to epileptic seizures, injuries, inflammatory processes; endocrine comas; gangrene and bedsores; severe immunodeficiency; underdevelopment or absence of organs in congenital malformations; prematurity; suicides in depressive states.

DECLARATIONS:

Disclosure Statement

The authors have no potential conflicts of interest to disclosure, including specific financial interests, relationships, and/or affiliations relevant to the subject matter or materials included.

Statement of Ethics

The authors have no ethical conflicts to disclosure.

Data Transparency

The data can be requested from the authors.

Funding Sources

There are no external sources of funding.

Consent for publication

All authors give their consent to publication.

References

1. Holovanova IA, Shevchenko AS. The issue of patient-oriented organization of palliative and hospice care in Ukraine. *Experimental and Clinical Medicine*. 2021;90(2):21-7. DOI: 10.35339/ekm.2021.90.2.hos.
2. Nesterenko VG. The controllability of pain in palliative medicine as a factor in the medical and social impact on the patients' quality of life at the end-of-life. *Experimental and Clinical Medicine*. 2024;93(1):69-86. DOI: 10.35339/ekm.2024.93.1.nes. [In Ukrainian].
3. Smiianov VA. Medical assistance in dying for palliative patients in different countries of the world: lessons on euthanasia legalization. *Inter Collegas*. 2024;11(2):32-40. DOI: 10.35339/ic.11.2.smi.
4. Nesterenko VG, Shevchenko AS, Zelenska KO, Hryhorov MM. Clinical and epidemiological characteristics of disabling neurological diseases (literature review). *International Neurological Journal*. 2024;20(4):176-84. DOI: 10.22141/2224-0713.20.4.2024.1079. [In Ukrainian].
5. Kluger BM, Hudson P, Hanson LC, Buzgova R, Creutzfeldt CJ, Gursahani R, et al. Palliative care to support the needs of adults with neurological disease. *Lancet Neurol*. 2023;22(7):619-31. DOI: 10.1016/S1474-4422(23)00129-1. Erratum in: *Lancet Neurol*. 2023;22(10):e11. DOI: 10.1016/S1474-4422(23)00310-1. PMID: 37353280.
6. Lekhan VM. Impact of different types of pain on organizational decisions in palliative medicine. *Medicine Today and Tomorrow*. 2024;93(1):47-55. DOI: 10.35339/msz.2024.93.1.lvm.
7. Nesterenko VG, Mykhnevych KG. Practical aspects of pain relief in palliative medicine. *Inter Collegas*. 2024;11(1):57-72. DOI: 10.35339/ic.11.1.nem.
8. Holovanova IA, Shevchenko AS. Determination of the main needs of palliative patients and ways of their provision in the health care system of Ukraine. *Experimental and Clinical Medicine*. 2021;90(3):88-94. DOI: 10.35339/ekm.2021.90.3.hos.
9. Lekhan VM. The importance of medical cannabis for solving the problem of providing analgesia to palliative patients in Ukraine. *Experimental and Clinical Medicine*. 2024;93(1):87-93. DOI: 10.35339/ekm.2024.93.1.lvm.
10. Wijdicks EF, Varelas PN, Gronseth GS, Greer DM; American Academy of Neurology. Evidence-based guideline update: determining brain death in adults: report of the Quality Standards Subcommittee of the American Academy of Neurology. *Neurology*. 2010;74(23):1911-8. DOI: 10.1212/WNL.0b013e3181e242a8. PMID: 20530327.
11. Shevchenko AS, Brown GW. What evidence-based medicine can oppose to falsifications in science. *Bull KhRIPHS*. 2020;96(4):24-35 DOI: 10.5281/zenodo.5076629. [In Ukrainian].
12. Nesterenko VG, Burzyńska J. Justification of the model of optimized system of providing palliative and hospice care to the population of Ukraine. *Inter Collegas*. 2024;11(3):10p. In press. DOI: 10.35339/ic.11.3.neb.
13. The need for palliative care: an assessment based on 2018 data. Ukrainian Center for Public Data. [Internet]. Available at: <https://socialdata.org.ua/palliative/> [accessed 20 Dec 2024]. [In Ukrainian].
14. Nesterenko VG. The need for palliative and hospice care in Ukraine in 2018–2020. *Medicine Today and Tomorrow*. 2021;90(3):43-52. DOI: 10.35339/msz.2021.90.3.nes. [In Ukrainian].
15. Nesterenko VG. Forecasting the need for palliative and hospice care in Ukraine for 2021–2022. *Experimental and Clinical Medicine*. 2021;90(4):25-34. DOI: 10.35339/ekm.2021.90.4.nes. [In Ukrainian].
16. Nesterenko VG, Redka IV, Sukhonosov RO, Grygorov SM, Shevchenko AS, Aliieva TD. Forecasting the need for palliative and hospice care using the creeping trend method with segment smoothing. *Wiad Lek*. 2024;77(5):980-4. DOI: 10.36740/WLek202405116. PMID: 39008586.
17. Nesterenko VG, Shevchenko VV. Calculation of the national need for palliative care by the refined method of trends in the conditions of a military crisis. *Eastern Ukrainian Medical Journal*. 2024;12(3):711-20. DOI: 10.21272/eumj.2024;12(3):711-720.
18. Nesterenko VH, Peresyphina TV, Shevchenko OS, Sarkhadova II. Palliative and hospice care in the Kharkiv region in the first year of the war. *Ukraine. Nation's Health*. 2023;73(3):106-12. DOI: 10.32782/2077-6594/2023.3/18.
19. Nesterenko VG, Sarkhadova II. Palliative and hospice care for adults in the Kharkiv Regional Center of Palliative Medicine "Hospice" in 2016–2022. *Experimental and Clinical Medicine*. 2023;92(2):58-68. DOI: 10.35339/ekm.2023.92.2.nes. [In Ukrainian].
20. Nesterenko VG. Peculiarities of using the SF-36 questionnaire in palliative neurological patients. *Medicine Today and Tomorrow*. 2023;92(2):62-72. DOI: 10.35339/msz.2023.92.2.nes. [In Ukrainian].

21. Nesterenko VG. Peculiarities of using the SF-36 questionnaire in palliative oncological patients. *Experimental and Clinical Medicine*. 2023;92(3):73-85. DOI: 10.35339/ekm.2023.92.3.nes. [In Ukrainian].
22. Nesterenko VG, Lytvynenko OYu. Readiness for dialogue regarding euthanasia of palliative patients in Ukraine. *Experimental and Clinical Medicine*. 2024;93(2):79-88. DOI: 10.35339/ekm.2024.93.2.nel. [In Ukrainian].
23. Titko EV, Deineko OV. Euthanasia / suicide tourism as the human rights protection phenomenon in the present time. *Academic notes of the V.I. Vernadsky Tavrida National University. Series: Legal Sciences*. 2020;31(70(2(3))):227-32. DOI: 10.32838/2707-0581/2020.2-3/36. [In Ukrainian].
24. Shevchenko AS, Kuts YuV, Gavrylova SA, Gavrylov EV. Legislative, historical and practical aspects of phenomena that potentially violate the right to life. *Bull KhRIPHS*. 2020;93(1):20-7. DOI: 10.5281/zenodo.3960410. [In Ukrainian].
25. Nesterenko VG, Krasnikova SO, Shevchenko AS. Media discussions about hospice and palliative care in Ukraine since the beginning of the war. Proceedings of the VI-th scientific and practical conference with international participation "Public health in Ukraine: problems and ways to solve them", dedicated to the 100-th anniversary of the department of public health and health care management of the Kharkiv National Medical University (Tomilin's readings), 2 Nov 2023. 4 p. DOI: 10.5281/zenodo.10053065.
26. Holovanova IA, Shevchenko AS. Evaluation of packaged funding programs for palliative and hospice care by the National Health Service of Ukraine. *Experimental and Clinical Medicine*. 2021;90(4):45-52. DOI: 10.35339/ekm.2021.90.4.hos.
27. Ortega CE, Seidner Y, Dominguez I. Mining CK2 in cancer. *PLoS One*. 2014;9(12):e115609. DOI: 10.1371/journal.pone.0115609. Erratum in: *PLoS One*. 2015;10(3):e0120224. DOI: 10.1371/journal.pone.0120224. PMID: 25541719.
28. Yao N, Sun X. Challenges and Opportunities to Conduct Cancer Care Research in China: Experience from a Pilot Project. *China Health Rev*. 2014;5(2):4-9. PMID: 25664325.
29. Mishchenko M, Shevchenko A, Mishchenko A. Key directions in the organization of health care and prevention actions to reduce mortality from cardiovascular diseases in Ukraine. Abstracts of XVI international scientific and practical conference "Prospects for the development of modern science and practice" (Graz, Austria, 11 May 2020). P. 80-2. DOI: 10.5281/zenodo.3820773.
30. Ognev VA, Mishchenko MM, Mishchenko AN, Shevchenko AS. World strategy on tobacco control and mortality from cardiovascular diseases in Ukraine. Abstracts of the 2nd International scientific and practical conference "Modern science: problems and innovations" (Stockholm, Sweden, 03–05 May 2020). SSPG Publish. P. 102-8. DOI: 10.5281/zenodo.3787953.
31. Mishchenko MM, Shevchenko AS, Mishchenko AN. Risk factors of ischemic stroke and their interconnection. Dynamics of the development of world science: Abstracts of the 8th International scientific and practical conference (Vancouver, Canada, 15-17 April 2020). Publishing House "ACCENT". P. 124-9. DOI: 10.5281/zenodo.3755806.
32. Ognev VA, Mishchenko MM, Mishchenko AN, Shevchenko AS. About value and study of the various risk factors for the development of a cerebral stroke. Topical issues of the development of modern science: Abstracts of the 8th International scientific and practical conference (Sofia, Bulgaria, 08-10 Apr 2020). Publishing House "ACCENT". P. 62-9. DOI: 10.5281/zenodo.3747558.
33. Mishchenko MM, Mishchenko AN, Shevchenko AS. On the study of risk factors for cardiovascular diseases. Proceedings of the XIX International Scientific and Practical Conference "Modern Science and its Impact on Humanity Development" (Ukraine, Kharkiv, 28 Feb 2020). *Bull KhRIPHS*. 2020;93(1):60-1. DOI: 10.5281/zenodo.4005972.
34. Picard G, Bier JC, Capron I, De Deyn PP, Deryck O, Engelborghs S, et al. Dementia, End of Life, and Euthanasia: A Survey Among Dementia Specialists Organized by the Belgian Dementia Council. *J Alzheimers Dis*. 2019;69(4):989-1001. DOI: 10.3233/JAD-181277. PMID: 31127774.
35. Enroth L, Raitanen J, Hervonen A, Nosraty L, Jylha M. Is socioeconomic status a predictor of mortality in nonagenarians? The vitality 90+ study. *Age Ageing*. 2015;44(1):123-9. DOI: 10.1093/ageing/afu092. PMID: 25002455.
36. Fairthorne J, Hammond G, Bourke J, Jacoby P, Leonard H. Early mortality and primary causes of death in mothers of children with intellectual disability or autism spectrum disorder: a retrospective cohort study. *PLoS One*. 2014;9(12):e113430. DOI: 10.1371/journal.pone.0113430. PMID: 25535971.

37. Holland JN, Schmidt AT. Static and Dynamic Factors Promoting Resilience following Traumatic Brain Injury: A Brief Review. *Neural Plast.* 2015;2015:902802. DOI: 10.1155/2015/902802. PMID: 26347352.
38. Bely SN, Yeskova LI, Shevchenko AS. HIV: the history of detection, tropism and methods of immunization. *Red Ribbon: Journal for medical workers and AIDS service organizations.* 2000;(2):15-9. DOI: 10.5281/zenodo.3715389.
39. Cambier CJ, Falkow S, Ramakrishnan L. Host evasion and exploitation schemes of *Mycobacterium tuberculosis*. *Cell.* 2014;159(7):1497-509. DOI: 10.1016/j.cell.2014.11.024. PMID: 25525872.
40. Field N, Lim MS, Murray J, Dowdeswell RJ, Glynn JR, Sonnenberg P. Timing, rates, and causes of death in a large South African tuberculosis programme. *BMC Infect Dis.* 2014;14:3858. DOI: 10.1186/s12879-014-0679-9. PMID: 25528248.
41. Belayneh M, Giday K, Lemma H. Treatment outcome of human immunodeficiency virus and tuberculosis co-infected patients in public hospitals of eastern and southern zone of Tigray region, Ethiopia. *Braz J Infect Dis.* 2015;19(1):47-51. DOI: 10.1016/j.bjid.2014.09.002. PMID: 25467923.
42. Das S, Tucker I, Stewart P. Inhaled dry powder formulations for treating tuberculosis. *Curr Drug Deliv.* 2015;12(1):26-39. DOI: 10.2174/1567201811666140716123050. PMID: 25030114.
43. Salazar-Austin N, Dowdy DW, Chaisson RE, Golub JE. Seventy Years of Tuberculosis Prevention: Efficacy, Effectiveness, Toxicity, Durability, and Duration. *Am J Epidemiol.* 2019;188(12):2078-2085. DOI: 10.1093/aje/kwz172. PMID: 31364692.
44. Jelinska A, Zajac M, Dadej A, Tomczak S, Geszke-Moritz M, Muszalska-Kolos I. Tuberculosis – Present Medication and Therapeutic Prospects. *Curr Med Chem.* 2020;27(4):630-56. PMID: 30457045. DOI: 10.2174/0929867325666181120100025.
45. Global tuberculosis report 2024. Geneva: World Health Organization; 2024. 69 p. Available at: <https://iris.who.int/bitstream/handle/10665/379339/9789240101531-eng.pdf>
46. Rijal A, Nielsen EE, Hemmingsen B, Neupane D, Gæde PH, Olsen MH, Jakobsen JC. Adding exercise to usual care in patients with hypertension, type 2 diabetes mellitus and/or cardiovascular disease: a protocol for a systematic review with meta-analysis and trial sequential analysis. *Syst Rev.* 2019;8(1):330. DOI: 10.1186/s13643-019-1233-z. PMID: 31847918.
47. Kim SU, Kim BK, Park JY, Kim DY, Ahn SH, Park YB, et al. Fibrosis-4 index at diagnosis can predict all-cause mortality in patients with rheumatoid arthritis: A retrospective monocentric study. *Mod Rheumatol.* 2020;30(1):70-7. DOI: 10.1080/14397595.2018.1558760. PMID: 30557057.
48. Avouac J, Degraeve R, Vergneault H, Combiere A, Wanono S, Boisson M, et al. Risk of liver fibrosis induced by methotrexate and other rheumatoid arthritis medications according to the Fibrosis-4 Index. *Clin Exp Rheumatol.* 2022;40(1):150-7. DOI: 10.55563/clinexprheumatol/usddjr. PMID: 33938795.
49. Uzun GS, Bulat B, Ayan G, Kılıç L, Kalyoncu U. Liver fibrosis in inflammatory arthritis patients treated with methotrexate and hydroxychloroquine: A FIB-4 index analysis. *Int J Rheum Dis.* 2024;27(10):e15390. DOI: 10.1111/1756-185X.15390. PMID: 39439367.
50. Cannon MD, Martin P, Carrion AF. Bacterial Infection in Patients with Cirrhosis: Don't Get Bugged to Death. *Dig Dis Sci.* 2020;65(1):31-7. DOI: 10.1007/s10620-019-05943-6. PMID: 31768880.
51. European Pregnancy and Paediatric HIV Cohort Collaboration (EPPICC) Study Group in EuroCoord. Prevalence and Clinical Outcomes of Poor Immune Response Despite Virologically Suppressive Antiretroviral Therapy Among Children and Adolescents with Human Immunodeficiency Virus in Europe and Thailand: Cohort Study. *Clin Infect Dis.* 2020;70(3):404-15. DOI: 10.1093/cid/ciz253. PMID: 30919882.
52. Hussein M, Dibaba B, Wondu Y. Factors Influencing Effectiveness of Peer Education Against Non-adherence to Antiretroviral Therapy Among Adult People Living with HIV/AIDS: A Pre-Post Intervention Study in Arsi Zone, Ethiopia. *HIV AIDS (Auckl).* 2020;12:1-7. DOI: 10.2147/HIV.S228330. PMID: 32021482.
53. Ogard-Repal A, Berg RC, Fossum M. Peer Support for People Living With HIV: A Scoping Review. *Health Promot Pract.* 2023;24(1):172-90. DOI: 10.1177/15248399211049824. PMID: 34693785.
54. Haque MA. Seasonal Incidence of Community-acquired Pneumonia: A Retrospective Study in a Tertiary Care Hospital in Kathmandu, Nepal. *Cureus.* 2019 Dec 18;11(12):e6417. DOI: 10.7759/cureus.6417. PMID: 31988818.
55. Allen JL. Airway function throughout the lifespan: Pediatric origins of adult respiratory disease. *Pediatr Investig.* 2019;3(4):236-44. DOI: 10.1002/ped4.12165. PMID: 32851329.

56. Lin WY, Verma VK, Lee MY, Lin HC, Lai CS. Prediction of 30-Day Readmission for COPD Patients Using Accelerometer-Based Activity Monitoring. *Sensors (Basel)*. 2019;20(1):217. DOI: 10.3390/s20010217. PMID: 31905995.
57. Ogata-Suetsugu S, Hamada N, Tsuda T, Takata S, Kitasato Y, Inoue N, et al.; Fukuoka tobacco-related lung disease (FOLD) registry study group, Fukuoka, Japan. Characteristics of tobacco-related lung diseases in Fukuoka Prefecture, Japan: A prospective, multi-institutional, observational study. *Respir Investig*. 2020;58(1):74-80. DOI: 10.1016/j.resinv.2019.10.002. PMID: 31727581.
58. Provenzano M, Coppolino G, Faga T, Garofalo C, Serra R, Andreucci M. Epidemiology of cardiovascular risk in chronic kidney disease patients: the real silent killer. *Rev Cardiovasc Med*. 2019 Dec;20(4):209-20. DOI: 10.31083/j.rcm.2019.04.548. PMID: 31912712.
59. Navaneethan SD, Schold JD, Arrigain S, Kirwan JP, Nally JV Jr. Body mass index and causes of death in chronic kidney disease. *Kidney Int*. 2016;89(3):675-82. doi: 10.1016/j.kint.2015.12.002. PMID: 26880461.
60. Bhandari SK, Zhou H, Shaw SF, Shi J, Tilluckdharry NS, Rhee CM, et al. Causes of Death in End-Stage Kidney Disease: Comparison between the United States Renal Data System and a Large Integrated Health Care System. *Am J Nephrol*. 2022;53(1):32-40. DOI: 10.1159/000520466. PMID: 35016183.
61. Aune D, Sun X, Nie J, Huang W, Liao B, Wang Y. Self-reported chronic kidney disease and the risk of all-cause and cause-specific mortality: outcome-wide association study of 54 causes of death in the National Health Interview Survey. *BMC Nephrol*. 2022 Apr 30;23(1):165. DOI: 10.1186/s12882-022-02771-1. PMID: 35488232.
62. Banshodani M, Kawanishi H, Moriishi M, Shintaku S, Tsuchiya S. Association between Dialysis Modality and Cardiovascular Diseases: A Comparison between Peritoneal Dialysis and Hemodialysis. *Blood Purif*. 2020;49(3):302-9. DOI: 10.1159/000504040. PMID: 31851981.
63. Ramos AM, Fernandez-Fernandez B, Perez-Gomez MV, Carriazo Julio SM, Sanchez-Nino MD, Sanz A, et al. Design and optimization strategies for the development of new drugs that treat chronic kidney disease. *Expert Opin Drug Discov*. 2020;15(1):101-115. DOI: 10.1080/17460441.2020.1690450. PMID: 31736379.
64. Martin C, Hamard C, Kanaan R, Boussaud V, Grenet D, Abely M, et al. Causes of death in French cystic fibrosis patients: The need for improvement in transplantation referral strategies! *J Cyst Fibros*. 2016;15(2):204-12. DOI: 10.1016/j.jcf.2015.09.002. PMID: 26391389.
65. Zang X, Monge ME, Gaul DA, McCarty NA, Stecenko A, Fernandez FM. Early Detection of Cystic Fibrosis Acute Pulmonary Exacerbations by Exhaled Breath Condensate Metabolomics. *J Proteome Res*. 2020;19(1):144-52. DOI: 10.1021/acs.jproteome.9b00443. PMID: 31621328.
66. Rosser BA, Chan C, Hoschitzky A. Surgical Management of Valvular Heart Disease in Mucopolysaccharidoses: A Review of Literature. *Biomedicines*. 2022;10(2):375. DOI: 10.3390/biomedicines10020375. PMID: 35203584.
67. Thomas S, Ramaswami U, Cleary M, Yaqub M, Raebel EM. Gastrointestinal Manifestations in Mucopolysaccharidosis Type III: Review of Death Certificates and the Literature. *J Clin Med*. 2021;10(19):4445. DOI: 10.3390/jcm10194445. PMID: 34640463.
68. Eskedal LT, Hagemo PS, Eskild A, Frøslie KF, Seiler S, Thaulow E. A population-based study relevant to seasonal variations in causes of death in children undergoing surgery for congenital cardiac malformations. *Cardiol Young*. 2007;17(4):423-31. DOI: 10.1017/S1047951107000881. PMID: 17583596.
69. Dodge OG, McKinney B. Causes of perinatal death in uganda africans, with special reference to congenital malformations. *J Pediatr*. 1964;65:267-9. DOI: 10.1016/s0022-3476(64)80529-1. PMID: 14198801.
70. Agrawal A, Swaminathan S, Guvvala SL, Gorla SR. Border Medicine: The Pediatric Cardiology Perspective. *Pediatr Cardiol*. 2020;41(1):202-5. DOI: 10.1007/s00246-019-02171-7. PMID: 31342117.
71. Szot JO, Campagnolo C, Cao Y, Iyer KR, Cuny H, Drysdale T, et al. Bi-allelic Mutations in NADSYN1 Cause Multiple Organ Defects and Expand the Genotypic Spectrum of Congenital NAD Deficiency Disorders. *Am J Hum Genet*. 2020 Jan 2;106(1):129-136. DOI: 10.1016/j.ajhg.2019.12.006. PMID: 31883644.
72. Almlı LM, Ely DM, Ailes EC, Abouk R, Grosse SD, Isenburg JL, et al. Infant Mortality Attributable to Birth Defects – United States, 2003–2017. *MMWR Morb Mortal Wkly Rep*. 2020;69(2):25-9. DOI: 10.15585/mmwr.mm6902a1. PMID: 31945037.
73. Dangel T, Wojciechowska U. New catalogue of conditions that may require pediatric palliative care. *Przegl Epidemiol*. 2020;74(3):503-20. DOI: 10.32394/PE.74.44. PMID: 33576589.

74. Lemmon ME, Bonifacio SL, Shellhaas RA, Wusthoff CJ, Greenberg RG, Soul JS, et al.; Neonatal Seizure Registry. Characterization of Death in Infants with Neonatal Seizures. *Pediatr Neurol.* 2020;113:21-5. DOI: 10.1016/j.pediatrneurol.2020.08.002. PMID: 32980743.
75. Barry A, Prentice T, Wilkinson D. End-of-life care over four decades in a quaternary neonatal intensive care unit. *J Paediatr Child Health.* 2023;59(2):341-5. DOI: 10.1111/jpc.16296. PMID: 36495233.
76. Reddihough DS, Baikie G, Walstab JE. Cerebral palsy in Victoria, Australia: mortality and causes of death. *J Paediatr Child Health.* 2001;37(2):183-6. DOI: 10.1046/j.1440-1754.2001.00644.x. PMID: 11328476.
77. Peterson MD, Wilson AM, Hurvitz EA. Underlying Causes of Death among Adults with Cerebral Palsy. *J Clin Med.* 2022;11(21):6333. DOI: 10.3390/jcm11216333. PMID: 36362560.
78. Durufle-Tapin A, Colin A, Nicolas B, Lebreton C, Dauvergne F, Gallien P. Analysis of the medical causes of death in cerebral palsy. *Ann Phys Rehabil Med.* 2014;57(1):24-37. DOI: 10.1016/j.rehab.2013.11.002. PMID: 24341986.
79. Alotaibi AS, Mahroos RA, Al Yateem SS, Menezes RG. Central Nervous System Causes of Sudden Unexpected Death: A Comprehensive Review. *Cureus.* 2022;14(1):e20944. DOI: 10.7759/cureus.20944. PMID: 35004089.
80. Ekkert A, Saulyte M, Jatuzis D. Inflammatory Disorders of the Central Nervous System Vessels: Narrative Review. *Medicina (Kaunas).* 2022;58(10):1446. DOI: 10.3390/medicina58101446. PMID: 36295606.
81. Nageshwaran S, Wilson HC, Dickenson A, Ledingham D. (Eds.). Inflammatory disorders of the central nervous system. P. 77-94. In: *Drugs in Neurology.* Oxford: Oxford Academic; 2016. DOI: 10.1093/med/9780199664368.003.0006.
82. Beghi E. The Epidemiology of Epilepsy. *Neuroepidemiology.* 2020;54(2):185-91. DOI: 10.1159/000503831. PMID: 31852003.
83. Auzmendi J, Puchulu MB, Rodriguez JCG, Balaszczuk AM, Lazarowski A, Merelli A. EPO and EPO-Receptor System as Potential Actionable Mechanism for the Protection of Brain and Heart in Refractory Epilepsy and SUDEP. *Curr Pharm Des.* 2020;26(12):1356-64. DOI: 10.2174/1381612826666200219095548. PMID: 32072891.
84. Sumelahti ML, Hakama M, Elovaara I, Pukkala E. Causes of death among patients with multiple sclerosis. *Mult Scler.* 2010;16(12):1437-42. DOI: 10.1177/1352458510379244. PMID: 20826526.
85. Cutter GR, Zimmerman J, Salter AR, Knappertz V, Suarez G, Waterbor J, et al. Causes of death among persons with multiple sclerosis. *Mult Scler Relat Disord.* 2015 Sep;4(5):484-90. DOI: 10.1016/j.msard.2015.07.008. PMID: 26346799.
86. Karamyan A, Brandtner H, Grinzinger S, Chroust V, Bacher C, Otto F, et al. Causes of death in critically ill multiple sclerosis patients. *Acta Neurol Scand.* 2017;136(4):305-9. DOI: 10.1111/ane.12724. PMID: 28008594.
87. Chou IJ, Kuo CF, Tanasescu R, Tench CR, Tiley CG, Constantinescu CS, Whitehouse WP. Comorbidity in multiple sclerosis: its temporal relationships with disease onset and dose effect on mortality. *Eur J Neurol.* 2020;27(1):105-12. DOI: 10.1111/ene.14040. PMID: 31309645.
88. Shevchenko AS, Shevchenko VV, Brown GW. The preventive direction of modern theories of health and health-saving in public health and education. *Inter Collegas.* 2024;11(1):45-51. DOI: 10.35339/ic.11.1.ssb.
89. Alieva TD, Shevchenko AS. On the issue of reproductive losses prevention in Ukraine. *Inter Collegas.* 2021;8(1):59-66. DOI: 10.35339/ic.8.1.59-66.
90. Shevchenko AS, Shevchenko VV, Shumskiy OL, Brown GW, Kucherenko SM, Kucherenko NS, Gavrylov EV. Motivational and regulatory instruments of valeological education. *Inter Collegas.* 2024;11(3):45-52. DOI: 10.35339/ic.11.3.sss.
91. Shevchenko A, Kucherenko S, Komyshan A, Shevchenko V, Kucherenko N. Formation of valeological competence in conditions of classroom and distance learning. *Scientific Notes of the Pedagogical Department.* 2022;(50):137-47. DOI: 10.26565/2074-8167-2022-50-14.
92. Smiianov VA, Zaitseva HS, Kurganskaya VA, Dyachenko AG, Zbarazhskiy VP, Smiianov YV, Pilipec OA. Vaccination coverage rates and the incidence of vaccine preventable diseases among children in sumy region of Ukraine. *Wiad Lek.* 2019;72(2):255-60. PMID: 30903783.

93. Golovanova IA, Smiiianov VA, Khorosh MV, Smiiianov YV, Smiiianova LI. Risk factors of smoking among students of higher medical educational establishments of Ukraine. *Wiad Lek.* 2018;71(6):1175-81. PMID: 30267496.

Received: 22 Oct 2024

Accepted: 31 Dec 2024

Cite in Vancouver style as: Smiiianov VA, Hubert-Lutecka A. Analytical study of the leading causes of death of palliative patients. *Inter Collegas.* 2024;11(4):10p. In press. <https://doi.org/10.35339/ic.11.4.smh>

Creative Commons license (BY-NC-SA) Smiiianov V.A., Hubert-Lutecka A., 2024