

PEDIATRICS

M.A Gonchar, E.V. Omelchenko, A.S. Senatorova, M.I. Strelkova, A.E. Silicheva, M.N. Ermolaev

THE STATE OF THE CARDIOVASCULAR SYSTEM IN CHILDREN WITH GASTROINTESTINAL DISORDERS

Kharkiv National Medical University, Ukraine

Summary. The article deals with the main approaches to stratification of cardiovascular risk in children with gastrointestinal disorders. The study focuses on unmodified risk factors identified by "Genetic questionnaire". Folate cycle gene polymorphism was shown to promote the development of cardiovascular diseases in children.

Keywords: cardiovascular risk, children, cardiovascular system, gastrointestinal diseases.



INTRODUCTION

A rapid increase in development of preventive medicine based on measures taken to avoid occurrence of diseases or to diagnose and treat existent diseases at early stages has been observed recently. Numerous risk factors for cardiovascular diseases (CVD) are divided into modified and unmodified ones [1]. Modified risk factors include such factors as smoking, dyslipidemia (increased LDL, triglycerides, reduced HDL), high blood pressure, diabetes, obesity, dietary factors, low level of physical activity, alcohol abuse. Unmodified risk factors are personal life history and family history [2].

Generally accepted criteria used to identify the severity of CVD risk implicate the so-called new lipid and nonlipid factors.

New lipid risk factors comprise elevated triglycerides, lipoprotein remnants, small particles of LDL, HDL subtypes, apolipoproteins B and A-I, cholesterol LDL/cholesterol HDL ratio [3].

New nonlipid risk factors include serum homocysteine; thrombogenic/antithrombogenic factors (platelets and clotting factors, fibrinogen, activated factor VII, plasminogen activation inhibitor-1 (PAI-1), tissue plasminogen activator (tPA), von Willebrand factor (vWF), V Leiden factor, protein C, antithrombin III); inflammatory factors; increased fasting glucose levels [4].

The study provided evidence for an important role of folate cycle gene polymorphism in the development of cardiovascular diseases. The aforementioned process can be caused by a disturbance in DNA synthesis resulting in dysregulation of proliferative processes and apoptosis [5]. Disturbances in folate cycle are thought to be caused by genetic defects of enzymes (MTHFR, MTR, MTRR), folic acid deficiency, vitamins B6 and B12 deficiency. The course of CVD is more complicated by a combination of several, even moderately expressed above-mentioned risk factors as compared to one most important risk factor.

Additional risk factors for disturbances in folate cycle include gastrointestinal diseases with malabsorption of vitamin B complex (ulcerative colitis, Crohn's disease, celiac disease, enteritis, gastritis, peptic ulcer disease); malignant neoplasms of the pancreas and intestine; kidney diseases; persistent chronic infections; prolonged use of anti-

• **Corresponding Author:**

Olena Omelchenko, MD, PhD, Associate Professor, Department of Pediatrics 1 and Neonatology, Kharkiv National Medical University, Ukraine. E-mail: helenomelchenko@mail.ru

convulsants, methotrexate, metformin, H2-receptor antagonists, eufillin, folic acid antagonists, drugs interrupting the absorption of folates; factors related to lifestyle (smoking, excessive consumption of alcohol or coffee (more than 5 cups of coffee a day); psycho-emotional stress; sedentary way of life [6, 7].

Defects of the folate cycle enzymes are accompanied by microangiopathy of various focalization, particularly "bright" hands and feet, cutis marmorata; varix dilatation. Complications associated with homocysteinuria include mental retardation, mental disorders, convulsions, skeletal disorders, osteoporosis, dislocation of the lens, myopia, iridodonesis. CNS disorders caused by brain vessels thrombosis are manifested by spastic paralysis, paresis, mental retardation and neuro-psychic signs (poor attention switching, low performance capability). Challenges presented by risk stratification are rather associated with thorough clinical examination than with scales and classifications, as it is appropriate to assess the risk for each patient and adjust therapeutic approach depending on the findings.

2 PURPOSES, SUBJECTS AND METHODS:

2.1 Purpose

The aim of the study was to evaluate risk stratification factors for cardiovascular diseases in children with gastrointestinal disorders.

2.2 Subjects

The study was performed at Gastroenterology Department of Regional Children's Clinical Hospital in Kharkov from 2015 to 2016. 66 children aged from 2 to 17 years, 32 (48.5%) girls and 34 (51.5%) boys aged 11.3 ± 4.1 years with chronic gastrointestinal problems were examined. (chronic gastroduodenitis, duodenal ulcer, functional disorders of the biliary tract, pancreatopathy)

2.3 Methods

Diagnosis was verified by anthropometric data assessment, conventional clinical laboratory and instrumental examination (ECG, echocardiography). The original "Genetic questionnaire" was elaborated and used during the study.

Conflict of interests

There is no conflict of interests.

3 RESULTS AND DISCUSSION

Anthropometric data analysis showed normal BMI in 42.4% of the examined patients, increased BMI in 30.2% and low BMI in 24.0%. Thus, the group under investigation comprised children predisposed to obesity as well as patients with low body weight, which can be explained by loss of appetite, one of the symptoms of gastrointestinal diseases. The most prevalent gastrointestinal diseases were 42.4% chronic gastroduodenitis, 15.2% biliary dyskinesy, 12.1% pancreatopathy and 9.1% duodenal ulcer, which corresponds to the data provided by the leading experts concerning the current situation in pediatric digestive disorders in Ukraine.

At the time of admission all the children underwent objective evaluation of the cardiovascular system (table 1).

Table 1
Data on objective evaluation of the cardiovascular system in patients with gastrointestinal diseases

| Parameters | Data |
|-----------------------------------|-------------------|
| Average SBP, mmHg | 108.4 ± 11.05 |
| Average DBP, mmHg | 70.1 ± 7.8 |
| Average heart rate, beats per min | 82.2 ± 11.0 |

Thus, the analysis showed that 46 (69.7%) children had normal blood pressure, 7 (10.6%) children had prehypertension, and 9 (13.6%) children were found to have stage 1 hypertension. Meanwhile average systolic blood pressure was within a reference range. Muffled heart sounds were observed in 15 (22.7%) patients, functional systolic murmur in 25 (37.8%).

Only in 31.8% of patients ECG indices corresponded to age norms. The most frequent alterations included lengthening of the PQ interval, T wave and ST interval changes, lengthening of the QT interval; that is, patients were predominantly found to have ECG changes reflecting disturbances of myocardial repolarization (table 2). Evaluation of pacemaker function shown by electrocardiography revealed monotopic heart rhythm disturbances in 50% of patients.

Table 2

Changes of ECG indices in patients

| Character of changes in ECG indices | % of the number of children |
|--|-----------------------------|
| Lengthening of the PQ interval | 57.6 |
| Lengthening of the QRS interval | 18.2 |
| Lengthening of the QT interval | 24.2 |
| Changing of the ST interval and T wave | 42.4 |
| Monotopic heart rhythm disturbances: | 50.0 |
| – sinus arrhythmia | 21.2 |
| – sinus tachyarrhythmia | 10.6 |
| – sinus bradyarrhythmia | 18.2 |

Thus, sinus arrhythmia observed in 21.21% of patients was the most frequent condition. According to ECG data, no organic heart disease in any of the patients was registered which was also confirmed by echocardiography.

Minor structural cardiac abnormalities included abnormal trabeculae of the left ventricle in 28.8% and mitral valve prolapse (mainly grade 1) in 24.5% (table 3).

Table 3

The incidence of minor structural cardiac abnormalities in pediatric patients

| Anatomical and physiological indices | Number of patients (in %) |
|---|----------------------------|
| Mitral valve prolapse | 24.5% |
| Abnormal chords of the left ventricle | 28.8% |
| Hypertrophy of the papillary muscles and increased trabeculae apparatus of the left ventricle | 12.1% |

Echocardiography revealed moderate dilatation of the left ventricle in 30.3% (in patients with ulcerative colitis, Crohn's disease, celiac disease), other echocardiographic indices were normal.

The patients' medical records were studied in details to identify risk factors for CVS diseases.

Evaluation of patients' history using "Genetic questionnaire" showed that 37.9% of the patients' parents consumed more than five cups of coffee per day, 41.0% of parents smoke. Besides, 22.7% of parents were overweight, 11.4% suffered from class 1 obesity, 3.8% of parents had class 2 obesity. Type II diabetes was observed in 8.2% of parents and 18.2% other relatives. Furthermore, 36.4% of the surveyed children had sedentary lifestyle.

Thus, children with gastrointestinal disorders were found to have the following risk factors: family history burdened with early cardiovascular diseases in 41.1% of the examined children (7.6% of the respondents were diagnosed with cardiovascular diseases), 41.0% of smoking parents, overweight or obese 30.2% of children and 37.9% of parents.

The examined patients with gastrointestinal disorders suffered from the following cardiovascular abnormalities: prehypertension in 10.6%, stage 1 hypertension in 13.6%, muffled heart tones in 22.7%, functional systolic murmur in 37.8%.

Echocardiography detected deviations in 68.2% of the surveyed children; pacemaker function assessment showed monotopic heart rhythm disturbances in 50% of patients. ECG data confirmed by ECHO CG did not identify symptoms of organic heart diseases in any of the patients. Minor structural cardiac abnormalities included abnormal chordae of the left ventricle in 28.8% and mitral valve prolapse in 24.5%. Moderate dilatation of the left ventricle was observed in 30.3% of children.

The findings, obtained by "Genetic questionnaire" revealed phenotypic changes indicative of genetic defects of folate cycle enzymes, denoted by blonde hair in 54.5% of children, blue eyes - 28.8%, pale skin - 19.7%, vascular pattern on the skin - 13.6%, visual disorders - 13.6%, hearing disorders - 1.5%, posture abnormalities - 45.5%, predisposition to fracture of bones - 15.2% of the surveyed children (table 4). These changes can indicate the presence of microangiopathies of various localization if hyperhomocysteinemia.

Table 4
Findings obtained by "Genetic questionnaire" (%)

| Sign | Children | Parents | Siblings |
|---------------------------------------|----------|---------|----------|
| Blond hair | 54.5% | 68.5% | 53.0% |
| Blue eyes | 28.8 | 30.3% | 53.0% |
| Pale skin | 19.7% | 24.2% | 15.6% |
| Vascular pattern on the skin | 13.6% | 10.6% | 15.6% |
| Visual disorders | 13.6% | 20.0% | 18.8% |
| Hearing disorders | 1.5% | 1.5% | 13.6% |
| Diseases of the cardiovascular system | 7.6% | 28.0% | 3.2% |
| Posture abnormalities | 45.5% | 6.9% | 10.6% |
| Predisposition to fracture of bones | 15.2% | 3.1% | 1.5% |

Recognition of possible heterogeneity of these diseases provides basis for implication of the results of genetic studies in the selection and personalization of therapy and stratification of cardiovascular risk.

4 CONCLUSION

Determination of the state of the cardiovascular system in children with gastrointestinal disorders using "Genetic questionnaire" gives a possibility to stratify risk factors for the development of cardiovascular diseases.

REFERENCES

1. Conroy, R. M., Pyörälä K., Fitzgerald, A.P. et al. Estimation of ten-year risk of fatal cardiovascular disease in Europe: the SCORE project // *Eur. Heart J.* - 2009. - № 24 (11). - 987-1003.
2. Kathiresan S. Genetics of human cardiovascular disease // *Cell.* - 2012. - Mar 16. - № 148(6). - P. 1242-57.
3. Mancia, G. Total cardiovascular risk: a new treatment concept // *J. Hypertens. Suppl.* - 2015. - № 24 (2). - P. 17-24.
4. Willerson, J. T., Ridker P.M. Inflammation as a cardiovascular riskfactor // *Circulation.* - 2014. - № 109 (21, Suppl. 1). - 112-110.

5. Verdecchia, P., Angeli F., Pittavini L. et al. Regression of leftventricular hypertrophy and cardiovascular risk changes in hypertensive patients // *Ital. Heart J.* - 2012. - № 5 (7). - 505-510.
6. Association between decreased vitamin levels and MTHFR, MTR and MTRR gene polymorphisms as determinants for elevated total homocysteine concentrations in pregnant women / P. R. Barbosa, S. P. Stabler, A. L. Machado [et al.] // *Eur. J. Clin. Nutr.* - 2011. - Vol. 62, № 8. - 1010-1021.
7. Genetic polymorphisms involved in folate metabolism and concentrations of methylmalonic acid and folate on plasma homocysteine and risk of coronary artery disease / M. Biselli, A. R. Guerzoni, M. F. de Godoy [et al.] // *J. Thromb. Thrombolysis.* - 2010. - Vol. 29, № 1. - 32-40.

РЕЗЮМЕ

Гончарь М.О., Омельченко О.В., Сенаторова Г.С.,
Стрелкова М.І., Сілічева А.Е., Єрмолаєв М.М.
СТАН СЕРЦЕВО-СУДИННОЇ СИСТЕМИ У ДІТЕЙ З ГАСТРОЕ-
НТЕРОЛОГІЧНОЮ ПАТОЛОГІЄЮ

Харківський національний медичний університет

В статті викладено основні підходи до стратифікації кардіоваскулярного ризику у дітей с гастроентерологічною патологією. Зроблено акцент на немодифікованих факторах ризику, виявлених за допомогою «Генетичного опитувальника». Визначена роль поліморфізму генів фолатного циклу у розвитку серцево-судинних захворювань у дітей.

Ключові слова: кардіоваскулярний ризик, діти, серцево-судинна система, гастроентерологічна патологія.

РЕЗЮМЕ

Гончарь М.О., Омельченко О.В., Сенаторова Г.С.,

Стрелкова М.І., СілічеваА.Е., Ермолаєв М.М.

СОСТОЯНИЕ СЕРДЕЧНО-СОСУДИСТОЙ СИСТЕМЫ У ДЕТЕЙ С ГАСТРОЭНТЕРОЛОГИЧЕСКОЙ ПАТОЛОГИЕЙ

Харьковский национальный медицинский университет

В статье изложены основные подходы к стратификации кардиоваскулярного риска у детей с гастроэнтерологической патологией. Сделан акцент на немодифицированных факторах риска, выявленных с помощью «генетической опросника». Определена роль полиморфизма генов фолатного цикла в развитии сердечно-сосудистых заболеваний у детей.

Ключевые слова: кардиоваскулярный риск, дети, сердечно-сосудистая система, гастроэнтерологическая патология.

Received: 13-Feb. - 2016

Accepted: 27-Mar. - 2016